

The Effects of Pollution Reduction on a Wild Trout Stream

Final Report: 2005 - 2008



Spring Run

Dumpling Run



April 2009 (Rev 12/31/2009)

Prepared by W. Neil Gillies, Cacapon Institute

TABLE OF CONTENTS

| | |
|--|----|
| Executive Summary | 3 |
| Introduction | 5 |
| Partners | 7 |
| Methods | 8 |
| Hatchery Upgrade | 9 |
| Results | 11 |
| Flow | 11 |
| Water quality | 11 |
| How did water quality vary over time? | 14 |
| How did loads of key parameters vary over time? | 16 |
| Benthic Macroinvertebrate & Periphyton Analysis | 18 |
| Fisherman Survey | 20 |
| DISCUSSION | 23 |
| Erosion Control Projects | 24 |
| Outreach | 24 |
| Priority Watershed Planning | 24 |
| Year Four expectations | 25 |
| Appendix 1. Water Quality Statistics by Year | 26 |
| Appendix 2. Laboratory Methods for Water Quality Parameters. | 32 |
| Appendix 3. WV Save Our Streams Macroinvertebrate Assessment July 2003 | 33 |
| Appendix 4. Assessing the Condition of the Macroinvertebrate Communities of Spring Run .. | 34 |

Executive Summary

Spring Run is a unique aquatic resource in the Potomac Highlands region of West Virginia. It is fed by the largest spring in the region, with discharge typically ranging from 3000-3500 gallons per minute. With a temperature of ~53 °F at the spring and a pH of ~8, aquatic conditions are ideal for trout and the aquatic insects they eat. Spring Run flows about two miles from the spring source to its confluence with South Mill Creek, which is about four miles from the South Branch of the Potomac River. Spring Run has no tributaries. Much of the stream is shallow, and does not provide the complex habitat that trout need - but that is not the case in a one mile section in the middle of the Run.

Since the early 1960's, landowner's have issued permits for fly fishing, catch-and-release on about one mile of Spring Run. Landowners and other interested parties have installed and maintained various structures to form pools and overhead cover that provide hiding and feeding habitat for trout. Spring Run is recognized as one of the best "wild" rainbow trout fisheries in West Virginia. Friends of Springs Run's Wild Trout, was formed in 1996 to restore structure to Spring Run following flooding in 1996.

Beginning in 2001, fishermen noted a decline in the fishery. The number of large trout (14" and above) decreased and trout in the 11-13" range also declined in abundance. Fly fishermen also reported declines in aquatic insects, especially the "sulfur" mayfly which emerged in late spring and small yellow stoneflies, often called "yellow sallies".

Spring Run is rich in nutrients, delivered largely in effluent from the Spring Run Trout Hatchery (SRH - a trout rearing facility) which is located about one-third mile upstream from the upper end of the fly fishing section and about one-fourth mile below the spring. WVDEP issued a citation for violation of the Spring Run Trout Hatchery NPDES permit in January 2004, specifically for discharging excess biochemical oxygen demand (BOD) and total suspended solids (TSS). WVDNR, which operates SRH, installed an effluent treatment system at the facility to meet their permit requirements. It became operational in June 2007.

The pending construction of an effluent treatment system at SRH provided an opportunity to conduct a before/after study of the effects of the treatment system on the chemistry and biology of Spring Run. With funding from the Chesapeake Bay Program, the Spring Run environmental study began in 2005. The design of this project included an upstream & downstream component in Spring Run, and a treatment & control component comparing Spring Run and nearby Dumpling Run. Both streams are spring fed and have their origins in the same limestone and sandstone geology. This approach allows both within stream and between stream comparisons. Two years of baseline data and nearly two years of post-treatment data were collected.

The study parameters were: field chemistries (pH, temperature, dissolved oxygen, conductivity); laboratory chemistries (total phosphorus, various species of nitrogen, TSS, and BOD5); and biological (benthic macroinvertebrates, fish – by WVDNR and fisherman catch records by section, and stream flow. Water samples were collected

Results

Chemistry. Water chemistries were collected monthly from April through September. **Samples were not collected on hatchery cleanout days** because the biosolids from aquaculture effluent are notoriously patchy. Despite the wealth of confounding variables and within parameter variability, some patterns are reasonably clear from the study's data. The spring source water for the two streams has similar pH, conductivity, dissolved oxygen, TSS, and phosphorus, and supports similar communities of benthic macroinvertebrates. Source water in Dumpling Run tends to have less nitrate, and total N than Spring Run, and higher BOD5. Conductivity and pH tend to increase or not change in a downstream direction in Dumpling Run, and tend to decrease in a downstream direction in Spring Run. Nutrients and TSS are generally similar in the two Dumpling Run sites, and tend to increase in a downstream direction in Spring Run, often dramatically.

Review of post-treatment water quality data indicates that installation of the SRH treatment facility did not change the water quality characteristics of Spring Run's water downstream of the plant on non-cleanout days.

Phosphorus in particular remained elevated. TSS and BOD5 in Spring Run remained lower, on non cleanout sampling days, than in Dumpling Run.

It is clear that the sampling protocol did not capture reductions in the pollutant plume that occurred during cleanout as a result of the new effluent treatment system. Data provided by WVDNR indicates that this process reduced pollutant loads related to cleanout by roughly 90%. For example, based on WVDNR data: TP concentrations in the effluent stream during cleanout fell from an average of 3.2 mg/L to 0.38 mg/L; BOD fell from 33.05 mg/L to 3.67 mg/L; and TSS fell from 57.78 mg/L to 2 mg/L.

The volume of sludge removed monthly by the Spring Run treatment system varied widely. In 2008, the only year with a full record at the time of this writing, the average across twelve months was 12,375 gallons of sludge removed monthly from the site for land application.

Reports from Spring Run residents and fishermen indicate the decanted fluid released from the cleanup process into Spring Run is turbid, and has a foul odor. During the decant process, the turbid and pungent plume extends through the length of the fly fishing section.

Benthic Macroinvertebrates. Benthic invertebrate samples were collected twice each year at all sites, in spring and autumn, according to the standard protocols in use by the WVDEP. Benthic macroinvertebrate communities in all sites were characterized by dominance of either amphipods or chironomidae midge larvae.

The chironomidae, commonly an indicator of organic pollution, were only dominant in Spring Run at the top of the managed fishing section. Their relative abundance fell during the post treatment period, perhaps a sign that reductions in discharge of effluent solids due to installation of the hatchery treatment system were having a positive impact on the benthic macroinvertebrate community in Spring Run. A related indicator is that the subjective condition of the stream bottom at that site was notably improved (less odor, less “greasy feel” to sediment, less entrained algae) when sampled in the fall of 2008 (Craddock & Gillies, Personal Communication).

The natural dominance of amphipods in limestone streams may well have masked underlying benthic macroinvertebrate diversity at all sites. Diversity was higher in the lower Dumpling run site than in the two point source impacted sites in Spring Run.

Fish. Continuing decline in the Spring Run fishery was apparent throughout the study period. Fisherman catch records indicated that rainbow trout increase in abundance and size from the lower to the upper end of the managed catch and release section of Spring Run. Fish surveys by WVDNR found the fish population near the middle of the managed section characterized by very low diversity, with extreme dominance by rainbow trout. A survey near the confluence of Spring Run with South Mill Creek found greater diversity, including several abundant non-trout species, but still distinctly lower diversity than was observed in a similar survey in 1978.

Education and Outreach

Education and outreach were important components of this project. Friends of Spring Run’s Wild Trout cooperated with project partners, in hosting annual one-day stream monitoring workshops on Spring Run. The benthic workshops brought together a diverse group of individuals ranging from students; fly-fisherman, environmental professional and community leaders to better understand freshwater ecology. A display with a comprehensive overview of the study was shown at several conferences, including 2005 WV Watershed Celebration Day and the Volunteer Monitoring portion of the Mid-Atlantic Conference held in Canaan Valley. In addition, local newspapers published articles about the program

Partners: Friends of Spring Run’s Wild Trout, WV Conservation Agency; WV Department of Agriculture; WV Division of Natural Resources; WV Department of Environmental Protection; Cacapon Institute; Chesapeake Bay Program; and Freshwater Institute

The Effects of Pollution Reduction on a Wild Trout Stream Final Report: 2008

Introduction

Spring Run is a unique aquatic resource in the Potomac Highlands region of West Virginia. Unlike many small headwater streams that tend to go dry, it is fed by the largest spring in the region, with discharge typically ranging from 3000-3500 gallons per minute. With a temperature of ~53 °F at the spring and a pH of ~8, aquatic conditions are ideal for trout and the aquatic insects they eat. Spring Run flows about two miles from the spring source to its confluence with South Mill Creek, which is about four miles from the South Branch of the Potomac River. Spring Run has no tributaries. Much of the stream is shallow, and does not provide the complex habitat that trout need - but that is not the case in a one mile section in the middle of the Run.

Since the early 1960's, landowner's have issued permits for fly fishing, catch-and-release on about one mile of Spring Run. Landowners and other interested parties have installed and maintained various structures to form pools and overhead cover that provide hiding and feeding habitat for trout. Spring Run is recognized as one of the best "wild" rainbow trout fisheries in West Virginia. Friends of Springs Run's Wild Trout, was formed in 1996 to restore structure to Spring Run following flooding in 1996.

In the last few years, however, fishermen have noted a decline in the fishery. Emergence of the mayfly, Ephemerellidae (sulfurs) largely disappeared in the late 1990s. The number of large trout (14" and above) has decreased and trout in the 11-13" range have also declined in abundance. The population of trout is considerably lower in the lower reach of the one mile section. Algae formation is heavy in the upper reach of the catch-and-release section, much heavier than in the past, and algae reforms soon after washout by high water.

Spring Run is rich in nutrients, delivered largely in effluent from the Spring Run Trout Hatchery (SRH) which is located about one-third mile upstream from the upper end of the fly fishing section and about one-fourth mile below the spring. (SRH is a rearing facility; trout are not spawned there). In recent years, however, SRH has been producing more rainbow and "golden trout" for stocking West Virginia streams, and it seems that the effluent stream now may be a problem for the health of Spring Run. WVDEP issued a citation for violation of the Spring Run Trout Hatchery NPDES permit in January 2004, specifically for discharging excess biochemical oxygen demand (BOD) and total suspended solids (TSS). WVDNR, which operates SRH, installed an effluent treatment system at the facility to meet their permit requirements. It became operational in June 2007.

The pending construction of an effluent treatment system at SRH provided a unique opportunity to address a number of issues of both regional and national significance:

1. Will the hatchery effluent treatment process significantly reduce nutrient discharge? Fish hatcheries throughout the country produce nutrient-rich effluents of concern to receiving waters. This study will evaluate the downstream result of effluent reduction of BOD and TSS, as well as nutrients, from a small but high throughput point source. The results of renovation at SRH and this study will provide important information to the WV Potomac Tributary Strategy point source innovation process.
2. What are the biological impacts of Spring Run's high nutrient levels, and how is the biota affected by reductions in nutrients, TSS and BOD following hatchery upgrades? This issue is of importance to the nutrient criteria development process that WV and the other 49 states are currently struggling through, as one of the key questions is: "what does nutrient impairment look like?"


3. Is the wild trout population in Spring Run being harmed by hatchery effluent, and does improvement in that effluent improve the trout fishery?
4. Is the benthic invertebrate population in Spring Run being harmed by hatchery effluent, and does improvement in that effluent improve diversity? Spring Run fishermen have noted the loss in recent years of a certain family of mayflies, the Ephemerellidae (Spiny crawler mayfly) that used to emerge regularly in the springtime. Also, WV DEP's Tim Craddock completed a benthic assessment of Spring Run in 2002, and found the lower part of the fly fishing section to be dominated by Chironomidae (midge) larvae, a group often indicative of pollution by organic waste.
5. Why do trout, especially larger fish, favor the upper part of the fly-fishing section? Why has the density-center of the trout population moved upstream in recent years? Is there a relationship between distribution of benthic invertebrates in the stream and trout distribution? If the Ephemerellidae mayflies and other pollution sensitive macroinvertebrates rebound after the hatchery effluent is treated, will the trout population improve also? In particular, are trout avoiding areas they used to frequent that are now dominated by midge larvae? If upgrades to the hatchery reduce organics in the stream and also the midge populations, will trout return to those areas? If that turns out to be true, and we could demonstrate that it is true, that would buttress public acceptance of benthic invertebrate stream assessments.

Overall, this project has the potential to address many questions beyond the five questions identified above.

This progress report will provide an overview of two years of baseline data and the nearly two years of post-treatment data. Additional sampling of benthic macroinvertebrates and the fish community are planned, but unfunded at this time.

Timeline

The following figure provides an approximate timeline for significant events related to the fishery in Spring Run.



Partners


Friends of Spring Run's Wild Trout, Cacapon Institute (CI), the WV Conservation Agency (WVCA), WV Department of Agriculture (WVDA), WV Division of Natural Resources (WVDNR), WV Department of Environmental Protection (WVDEP), and the Freshwater Institute are partnering in this study. This project is funded primarily by West Virginia Conservation Agency's participation through the Chesapeake Bay Program. An associated erosion reduction project is funded through a Friends of Spring Run's Wild Trout 2005 Stream Partners Grant. Additionally, a home school group is monitoring the lower portion of Spring Run on a regular basis.

WVDA, WVDEP and WVDNR are all contributing in-kind services to the project. WVDA is collecting water samples, taking flow measurements, and performing field and laboratory water quality analyses. WVDEP is participating in collections of benthic invertebrate and periphyton and helping to cover the costs of analysis. WVDNR is performing fish surveys and Friends of Spring Run's Wild Trout is providing information on size and location of trout caught and released by permitted fly fisherman, and supplied equipment and labor for erosion-sediment control work..

The Freshwater Institute provided guidance to WVDNR on treatment methods for their effluent and is providing technical guidance for the project. WVCA is acting as project coordinator. Cacapon Institute has overall technical oversight for the project, will participate in field work, and will, in cooperation with partnering organizations, be responsible for data analysis and production of annual reports.

Methods

The project has two experimental components, an upstream/downstream design in Spring Run, and a control/experimental design that includes Dumpling Run, another spring fed stream nearby. Both streams have their origins in the same geology: limestone (Helderberg and Tonoloway/Wills Creek) and sandstone (Oriskany, McKenzie) formations. Spring Run flows off the ridge to the northwest into South Mill Creek, a tributary of the South Branch of the Potomac River. Dumpling Run flows east into the South Fork of the South Branch of the Potomac River.



The upstream/downstream part includes three sites in Spring Run: the first site is near the spring upstream of the hatchery; the second site is near the upper end of the fly fishing stream section; and the third is near the lower end of the fly fishing section. There are two sites on Dumpling Run, one just below the spring, the other some distance downstream. Overall, this design allows within stream and between stream comparisons. Under most conditions of flow the springs constitute the main source of water in both streams, but both streams also have periodic surface flow entering the main channel upstream of the spring. Due to unanticipated delays in construction of the effluent treatment system, the baseline period of data collection lasted for approximately two years (April 2005-May, 2007).

Water chemistries were collected monthly from April through September, typically on Wednesday. We chose to avoid collections on Mondays at the time of the pre-treatment hatchery cleanout because the "biosolids from the aquaculture effluent are notoriously patchy and difficult to characterize in sampling. . . . my thoughts on the nutrients is to focus on the residual chronic impacts, not the pulse of the cleaning plume"


(Joe Hankins, Freshwater Institute, personal communication). However, due to scheduling requirements, samples in September 2006 were collected on a Monday during the cleanout.

Water quality parameters include nitrogen in the forms of ammonia-nitrogen, nitrate/nitrite, total Kjeldahl nitrogen, total nitrogen (the sum of nitrate/nitrite and TKN), soluble reactive phosphorus, total phosphorus, total suspended solids (TSS), biochemical oxygen demand (BOD₅), and basic field parameters (pH, temperature, conductivity) (see Appendix 2 for laboratory methods). Flow measurements are collected at the same time as water samples at one site in each stream. This work is done primarily by the WVDA.

Benthic invertebrate and periphyton samples were collected twice each year at all sites, in spring and autumn, according to the standard protocols in use by the WVDEP. WVDEP format Rapid Bioassessment Protocol habitat analyses is conducted once each year. WVDEP and Cacapon Institute are primarily responsible for this fieldwork.

WVDNR will conduct electro shocking fishery assessments, and the permitted fly fishermen of Spring Run have been enlisted to record information on size and location of trout caught and released.

The methods used to analyze water quality data were graphical and statistical. Data distributions were displayed using box plots (sample box-plot at right), which are useful for side-by-side visual comparisons of data distributions. One way analysis of variance (ANOVA) was run on rank transformed data for comparison of median concentration distributions. An alpha value of 0.05 was used as the threshold for statistical significance. If a significant difference among group medians was detected, Tukey's multiple comparison test was used on the rank transformed data to determine where differences were located (Helsel and Hirsh, 1992). Statistics were calculated using JMP Statistical Discovery Software (version 4.0.2). Summary statistics and raw data are provided in Appendix XX.



Hatchery Upgrade (per Mike Shingleton, WVDNR)


The Spring Run Trout Hatchery effluent treatment system became fully operational on June 4, 2007. An automated composite sampler was installed as part of the hatchery renovation and was first used in April.

Cleaning process: Blocking weirs are placed in front of the quiescent zone of each raceway prior to cleaning; a standpipe is removed and the quiescent zone is brushed cleaned; 3-4 raceways are cleaned at a time; the wastewater from the quiescent zones is piped into the clarifier while water from raceways not being cleaned is discharged to Spring Run. The clarifier is filled to its holding capacity but is not allowed to overflow. Wastewater in the clarifier is allowed to settle 24-48 hours and the clarified water is then decanted and mixed with hatchery water back into Spring Run. The sludge remaining in the clarifier is pumped to the sludge holding tank for later disposal by land application (see table next page). The decanting process of discharging water from the clarifier occurs 2-3 times per week and lasts from 1 1/2 to 2 hours each time.

Private communications from Spring Run residents and fishermen report the decanted fluid released from the cleanup process is turbid, with a foul odor. The turbid decant plume extends through the length of the fly fishing section.

The following figures show reductions in total nitrogen, total phosphorus, BOD and TSS due to the treatment process. These results are based on composited samples collected during the cleanout process. Prior to the upgrade, these samples were collected during the actual cleanout. After the treatment system became operational, samples were collected during decanting.

Results of composite sample analysis of Spring Run Hatchery effluent during cleanout before and after treatment system became operational. The "before" period was from September 2006 through May 2007. The "after" period was from June through September 2007. Data provided by WVDNR.




| Spring Run Sludge Application | |
|---|-------------------|
| Data provided by Mike Shingleton, WVDNR | |
| Date | Gallons of Sludge |
| 08/13/07 | 19,500 |
| 10/01/07 | 12,000 |
| 11/8&9/07 | 24,000 |
| 2007 Total | 55,500 |
| 01/16/08 | 16,500 |
| 03/12/08 | 10,500 |
| 05/01/08 | 14,000 |
| 06/12/08 | 21,500 |
| 07/17/08 | 14,000 |
| 08/13/08 | 16,000 |
| 09/24/08 | 19,000 |
| 11/06/08 | 23,000 |
| 11/25/08 | 14,000 |
| 2008 Total | 148,500 |
| 01/20/09 | 24,500 |
| 03/03/09 | 21,500 |
| 03/04/09 | 3,000 |
| 04/16/09 | 24,500 |
| 05/21/09 | 24,000 |
| 07/02/09 | 10,500 |
| 07/09/09 | 21,000 |
| 08/25/09 | 31,000 |
| 2009 Total Through 8/25 | 160,000 |

The table at left provides a summary of land applications of sludge removed from the trout rearing facility. Prior to installation of the treatment system, this material would have been delivered to the stream during cleanout. No data on the dry weight of solids removed by this process is available.

Results

The effluent treatment system at the hatchery became operational in June 2007, leaving four months (June through September) of post-operational data in 2007 and a full six months in 2008. Appendix 1 provides water quality statistics by year. Appendix 4 provides detailed benthic invertebrate statistics.

Flow. Stream flows on sampling days for the most downstream sites in Spring Run and Dumpling Run are given in Figure 1. Flow in Spring Run was always at least twice as high as in Dumpling Run. Sampling day flows in 2005 were much more variable than in following years, and the streams were notably low between July and September in 2007, and between July and September 2008 – six of the ten months of post-treatment data. Due to the relationship between flow and the water quality parameters measured, these extended low flow conditions have an impact on the ability to draw statistical conclusions as to effects of the treatment system.



Water Quality. Median values for pH, conductivity and dissolved oxygen by site and by year are provided in Table 1. These data indicate the source water in each stream was very similar, and these median values varied narrowly across all sites and all years. Both streams are alkaline, with moderately high conductivity, and high dissolved oxygen levels. No difference between pre-treatment (2005 and 2006) and post-treatment (2007) periods was evident.

Table 1. Median pH, conductivity, and dissolved oxygen by site and year.

| Site | Median pH | | | | Median Conductivity | | | | Median Dissolved Oxygen (mg/L) | | | |
|----------------------|-----------|------|------|------|---------------------|------|------|------|--------------------------------|------|------|------|
| | 2005 | 2006 | 2007 | 2008 | 2005 | 2006 | 2007 | 2008 | 2005 | 2006 | 2007 | 2008 |
| Dumpling Run @Spring | 7.8 | 7.6 | 7.5 | 7.5 | 287 | 254 | 257 | 273 | 10.3 | 10.6 | 10.5 | 9.4 |
| Dumpling Run Bottom | 8.1 | 7.8 | 7.6 | 8.3 | 284 | 264 | 272 | 291 | 10.4 | 10.6 | 10.3 | 9.7 |
| Spring Run @Spring | 7.9 | 7.7 | 7.7 | 8.1 | 296 | 330 | 358 | 331 | 10.3 | 10.5 | 10.1 | 9.8 |
| Spring Run Middle | 7.8 | 7.7 | 7.5 | 7.9 | 255 | 255 | 254 | 255 | 10.3 | 10.4 | 10.3 | 9.6 |
| Spring Run Bottom | 7.5 | 7.6 | 7.5 | 7.9 | 248 | 253 | 255 | 256 | 10.6 | 10.7 | 10.4 | 9.6 |

Statistical comparisons: pH was statistically lower in Spring Run Bottom than Spring Run @Spring and Dumpling Run Bottom. Conductivity was higher in Spring Run @Spring than Spring Run Bottom. No difference between sites for dissolved oxygen.


Pre-Post Comparisons: pH was statistically lower at Dumpling Run @Spring during the Post treatment period; Dissolved oxygen was statistically lower in Spring Run @Spring during the POST period than all Spring Run sites during the PRE period.

Median values for total phosphorus (TP) and total nitrogen (TN) by site and year are provided in Table 2. Source water TP was similar in each stream, and did not increase in the downstream site in the control stream (Dumpling Run). Both sites below the hatchery in Spring Run (Spring Run Middle and Spring Run Bottom)


had significantly higher median TP than all other locations. No difference in TP between pre-treatment (2005 and 2006) and post-treatment (June 2007 through September 2008) periods was evident.

Table 2. Median total phosphorus (TP) and total nitrogen (TN) by site and year.

| Site | TP (mg/L) Median | | | | TN (mg/L) Median | | | |
|----------------------------|------------------|-------|-------|-------|------------------|------|------|------|
| | 2005 | 2006 | 2007 | 2008 | 2005 | 2006 | 2007 | 2008 |
| Dumpling Run @Spring = DR2 | 0.028 | 0.054 | 0.041 | 0.028 | 0.341 | 0.36 | 0.31 | 0.37 |
| Dumpling Run Bottom = DR1 | 0.026 | 0.044 | 0.038 | 0.031 | 0.364 | 0.36 | 0.34 | 0.59 |
| Spring Run @Spring = SR3 | 0.025 | 0.049 | 0.036 | 0.038 | 0.641 | 0.57 | 0.52 | 0.69 |
| Spring Run Middle = SR2 | 0.075 | 0.103 | 0.106 | 0.068 | 0.887 | 0.63 | 0.78 | 0.88 |
| Spring Run Bottom = SR1 | 0.087 | 0.103 | 0.085 | 0.070 | 0.877 | 0.73 | 0.75 | 0.96 |



SR1 and SR2 had significantly higher total phosphorus than all other sites. There was no difference between Pre and Post treatment periods.




DR1 Pre and Post had significantly lower TN than SR1 Post. No other significant differences were detected.

Median total nitrogen was consistently but not significantly higher in the Spring Run source water than in Dumpling Run. Median TN did not increase in the downstream direction in the control stream (Dumpling Run). Both sites below the hatchery in Spring Run (Spring Run Middle and Spring Run Bottom) had higher


(but not significantly) TN than control sites in Dumpling Run and the source water in the Spring Run spring. No difference in TN between pre-treatment and post-treatment periods was evident.

Table 3. Median BOD5 and TSS by site and year.

| Site | Median BOD5 (mg/L) | | | | Median TSS (mg/L) | | | |
|----------------------------|--------------------|-------|-------|-------|-------------------|------|------|-------|
| | 2005 | 2006 | 2007 | 2008 | 2005 | 2006 | 2007 | 2008 |
| Dumpling Run @Spring = DR2 | 1.54 | 1.4 | 0.68 | 0.9 | 4.5 | 1.15 | 2.5 | 2 |
| Dumpling Run Bottom = DR1 | 1.515 | 1.1 | 0.605 | 0.795 | 2.08 | 5.5 | 4.5 | 4.5 |
| Spring Run @Spring = SR3 | 0.985 | 0.645 | 0.415 | 0.465 | 1.58 | 2.58 | 3.5 | 1.575 |
| Spring Run Middle = SR2 | 0.91 | 0.76 | 0.53 | 0.735 | 5.5 | 5 | 3 | 2.5 |
| Spring Run Bottom = SR1 | 1.01 | 0.425 | 0.415 | 0.54 | 6.5 | 7 | 6 | 3.5 |



DR1-Pre significantly higher than all but DR2-Pre and SR2-Pre.



No significant differences.



Median values for BOD5 and TSS by site and year are provided in Table 3 and graphic comparisons in the associated graphs. Source water BOD5 was distinctly (but not significantly) higher in Dumpling Run than Spring Run. There was no marked change in BOD5 in the downstream direction in either stream. However,

median BOD5 decreased from Pre to Post periods at all sites, significantly only in DR1 and DR2. TSS was similar in the source water for the two streams, with data ranging broadly. Median TSS tended to increase slightly (but not significantly) in a downstream direction in both streams.

How did water quality vary over time?

The following four time-series bar graphs and associated text show how total N, total P, TSS and BOD5 concentrations varied during the study period to date. Also shown on each graph is the average of the flows at the two flow stations for each sampling period; this was done for the sake of graphic simplification, justified because these values were very strongly correlated ($r^2 = 0.94$).

Time series bar graphs of total nitrogen, total phosphorus concentrations at all permanent study sites.





Total nitrogen (TN) varied widely and, based on correlation analysis, generally tracked with flows at all sites. The highest levels at all sites were observed in August 2005 during a high water event, with the exception of SpringRunBottom in April 2008, which was high due to a very high TKN (2.74 mg/l). TN was usually higher in all Spring Run sites than Dumpling Run. Elevated TN at SR Middle and SR Bottom in September 2006 was probably due to sampling that occurred on hatchery cleanout day. However, sampling in April 2007 also occurred on a cleanout day, and a similar increase in TN at SR Middle and SR Bottom was not

observed. TN remained elevated, relative to other sites, in the two point source impacted Spring Run sites after the treatment system became operational in June 2007.

Total phosphorus (TP) varied widely over time at all sites and did not apparently vary with flow levels. However, the highest TP concentrations at all sites except SR Middle and Bottom were recorded during an active runoff event in April 2006. The highest TP concentration at SR Middle occurred when sampling occurred on a hatchery cleanout day in April 2007. The highest TP concentrations at SR Bottom occurred on September 2005, May and June 2006, for no apparent reason. Elevated TP downstream of the hatchery was evident at all flows at SR Middle and Bottom (with a notable exception in August 2006 at SR Bottom). TP concentrations at the two point source sites were often distinctly different in 2006; this was not the case in 2005, 2007, and 2008. Elevated TP at SR Middle and SR Bottom in September 2006 may have been due to sampling that occurred on hatchery cleanout day; however, TP was similarly high at these two sites in September 2005 when cleanout was not occurring. TP remained elevated, relative to other sites, in the two point source impacted Spring Run sites after the treatment system became operational in June 2007.

Time series bar graphs of total suspended solids, and biochemical oxygen demand concentrations at all permanent study sites.




TSS varied widely and very roughly tracked with flows at all sites. The highest TSS levels for all sites were observed in August '05 during a high water event, but were not notably high during an active runoff event in April 2006. TSS concentrations were more consistently elevated at DR Lower, SR Middle and SR Bottom in 2006 than 2005. Elevated TSS at SR Middle and SR Bottom in

September 2006 and April 2007 may have been due to sampling on hatchery cleanout day; however, TSS was also high at both spring sampling sites in September 2006. TSS was similarly elevated at all sites during the low water period from July through September 2007.


BOD5 varied substantially between sites. Correlation analysis indicated that BOD5 at Spring Run point source impacted sites tended to vary with flow, while patterns of BOD5 concentrations in non point sites had no apparent relationship to flow. BOD5 concentrations were notably low during the one active runoff event in April 2006, and notably high at all sites except SR Spring during a high water event in August 2005.

How did loads of key parameters vary over time? The following four time-series bar graphs and associated text show how total N, total P, TSS and BOD5 loads (in pounds per hour) varied at the two flow station sites during the baseline and the June-Sept 2007 post treatment sampling period.

Time series bar graphs of total nitrogen and total phosphorus loads (in pounds per hour) at the two flow sites.




Total nitrogen loads varied widely and generally tracked with flows at all sites. As with TN concentrations, SR consistently had the higher TN loads. The highest loads at both sites were delivered during the three highest water events.




Total phosphorus loads varied much more widely over time in DR than SR. As with TP concentrations, SR consistently had the higher TP loads.

Time series bar graphs of total suspended solids and biochemical oxygen demand loads (in pounds per

hour) at the two flow sites.



TSS loads varied widely and roughly tracked with flows at both sites. The highest loads were observed in August 2005 during a high water event. SR consistently had the higher TSS loads.




BOD5 loads varied substantially between the two sites and roughly varied with flows. The highest loads at both sites were observed in August 2005 during a high water event.


Benthic Macroinvertebrate Analysis

An assessment of Spring Run in 2003 by WVDEP (Tim Craddock, 2003) collected benthic invertebrate samples at sites near those chosen for the current study. The study found low diversity at the lower station, where the most abundant family was the Chironomidae, an indicator of organic pollution. It also found abundant Gammaridae amphipods at all sites. (See Appendix 3 for results, as well as a commentary of the challenge of assessing Spring Run macroinvertebrates by WVDEP's Tim Craddock.)


Benthic samples for this project were collected twice each year, in the spring and in the autumn, at all water quality sampling sites. The methods used were the standard collection, processing and identification protocols of WV DEP, with a 200 count subsample. Summary data is provided in Appendix 4. Observations during benthic field collections indicated abundance, often overwhelming abundance, of amphipods in both streams (Craddock and Gillies, personal observations). Amphipods are often extremely abundant in limestone spring fed streams, and their abundance renders many standard benthic invertebrate indices less meaningful in assessing this type of stream. Numerical results indicate that the most abundant organisms, at all study sites, were either Gammaridae (amphipods) or Chironomidae (midges). The result of this abundance, combined with use of the standard 200 count subsampling method, often means that the actual diversity of organisms in a sample is masked.

This section focuses on four indices that were chosen for their utility in this setting: Percent EPT (Ephemeroptera - mayfly, Plecoptera - stonefly, Trichoptera - caddisfly); percent dominance; percent Gammaridae (amphipods); and percent Chironomidae (non biting midges). Discussion of results are presented in the box below the graph for each index.






Percent dominance is common metric where high numbers typically indicate poor water quality. As noted above, however, such metrics are problematic in limestone spring fed streams where dominance by amphipods (or isopods) is normal. All sites had relatively high dominance throughout the study period, with all sites having at least one incidence of >75% dominance. DR@Spring always had % dominance greater than 50.



Percent Gammaridae is not a standard metric. It is used here in recognition that amphipods are commonly abundant in limestone spring fed streams, are certainly very abundant in all the study sites, and their abundance makes a number of standard metrics unreliable. % Gammaridae varied widely at all sites, with no universal pattern apparent. A quick comparison of the % Gammaridae graph to the % Dominance graph indicates that amphipods were, in most cases, the dominant organism at every site except SR Middle. It is notable that amphipods became proportionately much more important at SR Middle from Spring 2007 through Fall 2008 than previously.



Fisherman Survey

Anglers with permits to fly fish, catch-and-release were invited, by a notice posted at the Spring Run parking area, to report the date fished, species, length, and stream location of their catch. The fly-fishing, catch-and-release section of Spring Run extends for about one mile. This section was arbitrarily divided into 10 sections, marked at streamside; Numbered 0 thru 9, beginning with 0 at the downstream boundary and increasing upstream. Sections were not of equal length. Anglers fished wherever they chose. Fishing sessions ranged from less than an hour to several hours. Anglers reported on a card designed with stream sections vs. 6 length categories, in inches; 0-7, 8-10, 11-13, 14-16, 17-19, 20-up. This card was available from a box located convenient to the parking area and next to a locked box for depositing completed reports. The parking area was adjacent to stream section Number 4. A member of the monitoring team collected reports frequently and summarized data monthly. The purpose of the study was to acquire data on number, size, and location of Spring Run trout, not to evaluate angler success.

Anglers cooperated in collecting data with a participation rate estimated above 80% for sessions fished. Summary data presented below are for April through December in 2005, January through December in 2006, 2007, and 2008. The most heavily fished period is April through September. Data presented in the following tables are for rainbow trout. A small number of brown, brook and golden trout were reported.

| Table 6. Spring Run angler catch reports. | | | | | | | | | | | | |
|--|-----------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|----------|
| Spring Run Angler Catch Reports, Rainbow Trout: April thru Dec 2005 | | | | | | | | | | | | |
| 65 Anglers Reporting 230 Fishing Sessions | | | | | | | | | | | | |
| Length | Stream Section | | | | | | | | | | Total | % |
| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | | |
| 0-7 | 70 | 108 | 77 | 130 | 220 | 335 | 201 | 142 | 58 | 47 | 1388 | 37.5 |
| 8--10 | 22 | 35 | 26 | 72 | 146 | 221 | 191 | 217 | 203 | 162 | 1295 | 35 |
| 11--13 | 7 | 5 | 17 | 27 | 39 | 75 | 75 | 89 | 170 | 175 | 679 | 18.3 |
| 14--16 | | 1 | | 16 | 25 | 23 | 33 | 27 | 29 | 86 | 240 | 6.5 |
| 17--19 | | | | 1 | 5 | 4 | 9 | 7 | 10 | 24 | 60 | 1.6 |
| 20--up | | | 1 | | 1 | 4 | 7 | 10 | 6 | 13 | 42 | 1.1 |
| Total | 99 | 149 | 121 | 246 | 436 | 662 | 516 | 492 | 476 | 507 | 3704 | |
| % by Section | 2.7 | 4 | 3.3 | 6.6 | 11.8 | 17.9 | 13.9 | 13.3 | 12.9 | 13.7 | | |
| 16.1 rainbow trout/angler session | | | | | | | | | | | | |

| Spring Run Angler Catch Reports, Rainbow Trout: Jan thru Dec 2006 | | | | | | | | | | | | |
|--|-----------------------|-----------|-----------|------------|------------|------------|------------|------------|------------|------------|--------------|----------|
| 76 Anglers Reporting 232 Fishing Sessions | | | | | | | | | | | | |
| Length | Stream Section | | | | | | | | | | Total | % |
| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | | |
| 0-7 | 25 | 46 | 42 | 89 | 134 | 153 | 112 | 46 | 30 | 33 | 718 | 31.6 |
| 8--10 | 18 | 14 | 20 | 49 | 103 | 109 | 121 | 134 | 64 | 66 | 698 | 30.7 |
| 11--13 | 4 | 10 | 18 | 18 | 34 | 46 | 77 | 104 | 109 | 136 | 536 | 23.6 |
| 14--16 | | 4 | 4 | 8 | 9 | 18 | 31 | 42 | 43 | 92 | 251 | 11 |
| 17--19 | | 1 | 1 | 1 | 3 | 2 | 2 | 9 | 4 | 19 | 42 | 1.8 |
| 20--up | | | | 1 | 3 | 1 | | 1 | 1 | 8 | 15 | 0.7 |
| Total | 47 | 75 | 85 | 160 | 286 | 329 | 343 | 336 | 251 | 354 | 2272 | |
| % by Section | 2.1 | 3.3 | 3.7 | 7 | 12.6 | 14.5 | 15.1 | 14.8 | 11 | 15.6 | | |
| 9.8 rainbow trout/angler session | | | | | | | | | | | | |


| Spring Run Angler Catch Reports, Rainbow Trout: Jan thru Dec 2007 | | | | | | | | | | | | |
|--|-----------------------|-----------|-----------|------------|------------|------------|------------|------------|------------|------------|--------------|--------------|
| 59 Anglers Reporting 211 Fishing Sessions | | | | | | | | | | | | |
| Length | Stream Section | | | | | | | | | | Total | % |
| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | | |
| 0-7 | 20 | 54 | 30 | 51 | 61 | 89 | 51 | 37 | 27 | 22 | 442 | 29.3% |
| 8-10 | 8 | 19 | 18 | 35 | 65 | 88 | 38 | 36 | 29 | 48 | 384 | 25.4% |
| 11-13 | 3 | 9 | 8 | 15 | 65 | 62 | 63 | 53 | 36 | 76 | 390 | 25.8% |
| 14-16 | | | 5 | 8 | 30 | 27 | 36 | 54 | 29 | 55 | 244 | 16.2% |
| 17-19 | | 1 | | 2 | 7 | 3 | 2 | 4 | 2 | 14 | 35 | 2.3% |
| 20-up | | | | | | 2 | 2 | 2 | 1 | 7 | 14 | 0.9% |
| SUMS | 31 | 83 | 61 | 111 | 228 | 271 | 192 | 186 | 124 | 222 | 1509 | |
| % by Section | 2.1% | 5.5% | 4.0% | 7.4% | 15.1% | 18.0% | 12.7% | 12.3% | 8.2% | 14.7% | | |
| 7.2 rainbow trout/angler session | | | | | | | | | | | | |

| Spring Run Angler Catch Reports, Rainbow Trout: Jan thru Dec 2008 | | | | | | | | | | | | |
|---|----------------|-------------|-------------|-------------|--------------|--------------|-------------|--------------|-------------|--------------|-------------|-------|
| 45 Anglers Reporting 171 Fishing Sessions | | | | | | | | | | | | |
| Length | Stream Section | | | | | | | | | | Total | % |
| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | | |
| 0-7 | 24 | 64 | 42 | 62 | 69 | 118 | 68 | 87 | 48 | 72 | 654 | 47.2% |
| 8-10 | 15 | 51 | 31 | 37 | 63 | 63 | 48 | 47 | 37 | 86 | 478 | 34.5% |
| 11-13 | 1 | 11 | 8 | 10 | 22 | 17 | 7 | 13 | 33 | 43 | 165 | 11.9% |
| 14-16 | | 1 | 2 | 4 | 7 | 7 | 3 | 2 | 7 | 35 | 68 | 4.9% |
| 17-19 | | | 1 | | 1 | 1 | | | | 11 | 14 | 1.0% |
| 20-up | | | | | 2 | | | 2 | 1 | 3 | 8 | 0.6% |
| SUMS | 40 | 127 | 84 | 113 | 164 | 206 | 126 | 151 | 126 | 250 | 1387 | |
| % by Section | 2.9% | 9.2% | 6.1% | 8.1% | 11.8% | 14.9% | 9.1% | 10.9% | 9.1% | 18.0% | | |

8.1 rainbow trout/angler session

Sixty five anglers reported 230 fishing sessions in 2005, 76 anglers reported 232 fishing sessions in 2006, 59 anglers reported 210 fishing sessions in 2007, and 45 anglers reported 171 fishing sessions in 2008. The number of rainbow trout caught per session declined from 16.1 to 9.8 to 7.2 (in 2005, 2006, and 2007, respectively), and rose slightly to 8.1 per session in 2008. At the request of DNR, fishermen were asked to report time of each session fished in 2008, in order to determine catch per hour. For 2008 the catch rate was 2.32 rainbow trout per hour. Assuming fishermen in 2005, 2006 and 2007 fished the same length of time each fishing session as fishermen in 2008, hourly catch rates calculated for rainbow trout were: 2005 - 4.61/hr; 2006 - 2.81/hr; and in 2007 - 2.05/hr.

The total catch by stream section is summarized in the following graph. In each sampling year, the greatest numbers were caught in sections “4” through “9.” The catch in every section declined sharply from a high in 2005.



DISCUSSION

The two study streams are impacted by a variety of potential sources of pollution, some readily apparent and some not. The Spring Run watershed contains the trout rearing facility, which is a known source of BOD, TSS and nutrients, as well as a number of non point sources including poultry houses, residences, roads, and occasional cattle. The Dumpling Run watershed has no point sources, and apparently no poultry houses, but includes residences and small farms with livestock, as well as a dirt and gravel road. In addition, the source springs in both watersheds originate in limestone and sandstone strata and show rapid changes (turbidity, increase in flow) following heavy precipitation; this is indicative of solution channel connections through limestone at the surface of the ground.

Despite the wealth of confounding variables and within parameter variability, some patterns are reasonably clear from the above data. The spring source water for the two streams has similar pH, conductivity, dissolved oxygen, TSS, and phosphorus, and supports similar communities of benthic macroinvertebrates. Source water in Dumpling Run tends to have less nitrate, and total N than Spring Run, and higher BOD5. Conductivity and pH tend to increase or not change in a downstream direction in Dumpling Run, and tend to decrease in a downstream direction in Spring Run. Nutrients and TSS are generally similar in the two Dumpling Run sites, and tend to increase in a downstream direction in Spring Run, often dramatically.

Review of post-treatment water quality data indicates that installation of the SRH treatment facility did not change the water quality characteristics of Spring Run's water downstream of the plant on non-cleanout days. Phosphorus in particular remains elevated. TSS and BOD5 in Spring Run remain lower, on non cleanout days, than Dumpling Run. However, it is also clear that our sampling protocol did not capture reductions in the pollutant plume that may have occurred during cleanout as a result of the new effluent treatment system (see Hatchery Upgrade section above). Data provided by WVDNR indicates that this process reduces pollutant loads related to cleanout by roughly 90%. For example, based on WVDNR data: TP concentrations in the effluent stream during cleanout fell from an average of 3.2 mg/L to 0.38 mg/L; BOD fell from 33.05 mg/L to 3.67 mg/L and; TSS fell from 57.78 mg/L to 2 mg/L.

The volume of sludge removed monthly by the Spring Run treatment system varied widely (see table page 8). In 2008, the only year with a full record, the monthly average across twelve months was 12,375 gallons of sludge removed from the site for land application..

It is important to note that reports from Spring Run residents and fishermen indicate the decanted fluid released from the cleanup process into Spring Run is turbid, and has a foul odor. During the decant process, the turbid and pungent plume of decanted solution extends through the length of the fly fishing section.

Benthic macroinvertebrate communities in all sites are defined by dominance of either amphipods or chironomidae midge larvae. The natural dominance of amphipods in limestone streams may well have masked underlying benthic macroinvertebrate diversity. Diversity is higher in the lower Dumpling run site than in the two point source impacted sites in Spring Run.

The change in relative abundance of chironomidae in Spring Run at the top of the managed fishing section is, perhaps, a sign that large reductions in discharge of effluent solids due to installation of the hatchery treatment system is having a positive impact on the benthic macroinvertebrate community in Spring Run. A related indicator is that the subjective condition of the stream bottom at that site was notably improved (less odor, less "greasy feel" to sediment, less entrained algae) when last sampled in the fall of 2008 (Craddock & Gillies, Personal Communication).

Decline in the Spring Run fishery was apparent throughout the study period. While there were some signs of Rainbow Trout spawning and recruitment in late 2008 through early 2009, it is too early to know if any improvement can be expected.

Erosion Control Projects

Friends of Spring Runs Wild Trout, supported by a WV Stream Partners grant, partnered with the WVCA to reduce channel erosion and the resulting sedimentation problem in Spring Run. In 2006 WVCA supplied the design and heavy equipment to reshape and seed 210 feet of severely eroding roadside channel immediately above the spring source of Spring Run. In 2007 Friends of Spring Runs Wild Trout, with volunteered design, labor and equipment, installed three sediment basins and numerous sediment check dams and sections of rip-rap in three storm water runoff ditches, totaling about 600 feet, from developed areas. Sediment check dams of wood construction were placed in four woodland hollows which funnel run-off into Spring Run. Tree seedlings were planted in work areas in early 2008 and 2009.

Landowners remain concerned about sediment entering Spring Run from public roadway ditches. It is the hope of the working group and Friends of Spring Run's Wild Trout that this will become more of a priority for WV Division of Highways in the coming months.

Cattle which have been observed within the stream and along the unstable banks of Spring Run above the fly fishing section for years were removed near the end of 2007, and fencing installed to prevent cattle access in the future.

Outreach

A paper on the Spring Run project was presented at the 2007 Virginia/West Virginia Water Research Symposium in November 2007 by Cacapon Institute. This paper was coauthored by Neil Gillies (CI), Carla Hardy (WVCA), and Tim Craddock (WVDEP).

Priority Watershed Planning

The Mill Creek Watershed, of which Spring Run is a part, was listed by the WVDEP as impaired for excess fecal coliform bacteria in 1996. Following a Total Maximum Daily Load (TMDL) study, EPA called for a 37% reduction in fecal coliform loadings. A watershed based plan to implement the TMDL was submitted to EPA during the winter of 2008 and is now being implemented. Mill Creek is also a priority watershed for West Virginia's Potomac Tributary Strategy's Implementation Program, a Chesapeake Bay Program initiative.

A working group comprised of representatives from West Virginia Conservation Agency, West Virginia Department of Agriculture, Cacapon Institute, USDA Natural Resources Conservation Service and WVU Extension Service has worked to identify areas within the watershed that would benefit from BMP installation. A voluntary survey was mailed to all landowners within the entire watershed during the spring of 2007 requesting that they rank their environmental concerns. This information has been tallied and the working group has plans to bring educational programs that address these concerns to the local community as the project progresses.

The above steps all contribute to a process that will bring funds to the watershed for the purpose of improving water quality in the Mill Creek watershed. The funds will be used for projects such as feedlot relocations, installation of streambank fencing, alternative livestock water development, riparian buffers, failing septic upgrades, and wetland restoration. Funding for this project is anticipated in the summer of 2009.

Post Study Expectations

Benthic macroinvertebrate sampling will continue for at least one year to determine if apparent improvements in Spring Run below the hatchery are sustained over time. Fisherman will continue to report “catch and release” results for the foreseeable future, and WVDNR will conduct additional electro-shocking in the future, when possible.

Spring Run Fishermen Commentary

Jerry Burke and Carl Rettenberger have written detailed reports about Spring Run, describing changes in this fishery over the years, suggesting causes of the significant decline in the fishery, and identifying concerns which need to be addressed if this unique and valuable resource, Spring Run, is to recover as a great trout fishery and restore the community benefits it formerly supported. Their reports, briefly summarized below, are available as companion pieces to this project report.

Burke: Spring Run Recollections

Burke covered his experience and observations with Spring Run from 1966 when he met Harrison Shobe, the landowner who began the fly fishing catch-and-release project, until May 2008. It is intended that this document be periodically updated. Drawing on ~2000 days fishing, nearly 4000 hours of stream habitat restoration and improvement, interaction with several hundred Spring Run fishermen and fishery professionals from several organizations, Burke’s document touches on a variety of topics: Background; Electro-shocking; Physical Habitat; Fishing & Rainbow Trout; Other Trout and Non-Trout Species; Spawning; Loss of Macro-invertebrates; Declines in Trout Size and Number; Other Uses of Spring Run; Spring Run Monitoring Program; What is Cause or Causes of Trout Decline? (Poaching, Sediment, Spring Run Trout Hatchery; Two Chicken Production (broiler) Facilities, Dead Trout, Feeding Fish, Dwellings); Research; How Can the Spring Run Rainbow Trout Fishery be Restored to its Former Glory?

Rettenberger: The Decline of the Spring Run Fishery

Rettenberger reviews the physical and chemical characteristics of Spring Run, discusses the issues of sedimentation, parasites, toxins, predation, stocking, spawning, survival and mortality, and food sources. He concludes that:

The need for the Spring Run trout rearing facility is not being questioned. However, it must be operated and maintained in the most neighborly and environmentally friendly manner possible.

The writer does, or at least chooses to believe that the hatchery’s new effluent treatment facility, made operational in June of 2007, will reduce the amount of solids in the “*cleanout plumes*”. It is hoped that this reduction in sediments will result in a healthier environment for both the benthic community and the rainbow trout population.

Will this prove to be true, only time and the continuation of the macroinvertebrates and chemical analysis studies for a few more years will tell!

Appendix 1. Water Quality Statistics by Year.

| Site | Yr | Minimum | Median | Maximum | Mean | Std.Dev. |
|-------------------------|------|---------|--------|---------|-------|----------|
| Ammonia-N (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 0.003 | 0.007 | 0.082 | 0.025 | 0.033 |
| Dumpling Run Bottom | | 0.003 | 0.007 | 0.079 | 0.028 | 0.037 |
| Spring Run @Spring | | 0.003 | 0.012 | 0.915 | 0.167 | 0.367 |
| Spring Run Middle | | 0.051 | 0.093 | 0.214 | 0.107 | 0.063 |
| Spring Run Bottom | | 0.017 | 0.043 | 0.161 | 0.070 | 0.059 |
| Dumpling Run @Spring | 2006 | 0.008 | 0.008 | 0.008 | 0.008 | 0.000 |
| Dumpling Run Bottom | | 0.008 | 0.008 | 0.008 | 0.008 | 0.000 |
| Spring Run @Spring | | 0.008 | 0.008 | 0.008 | 0.008 | 0.000 |
| Spring Run Middle | | 0.008 | 0.046 | 0.102 | 0.048 | 0.034 |
| Spring Run Bottom | | 0.008 | 0.008 | 0.041 | 0.019 | 0.017 |
| Dumpling Run @Spring | 2007 | 0.003 | 0.003 | 0.003 | 0.003 | 0.000 |
| Dumpling Run Bottom | | 0.003 | 0.003 | 0.003 | 0.003 | 0.000 |
| Spring Run @Spring | | 0.003 | 0.003 | 0.026 | 0.007 | 0.009 |
| Spring Run Middle | | 0.029 | 0.048 | 0.283 | 0.090 | 0.099 |
| Spring Run Bottom | | 0.003 | 0.003 | 0.153 | 0.028 | 0.061 |
| Dumpling Run @Spring | 2008 | 0.003 | 0.003 | 0.013 | 0.006 | |
| Dumpling Run Bottom | | 0.003 | 0.003 | 0.039 | 0.011 | |
| Spring Run @Spring | | 0.003 | 0.003 | 0.008 | 0.004 | |
| Spring Run Middle | | 0.003 | 0.066 | 0.214 | 0.081 | |
| Spring Run Bottom | | 0.003 | 0.003 | 0.046 | 0.013 | |
| Nitrate-N (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 0.17 | 0.23 | 0.38 | 0.26 | 0.079 |
| Dumpling Run Bottom | | 0.19 | 0.26 | 0.50 | 0.31 | 0.117 |
| Spring Run @Spring | | 0.37 | 0.48 | 0.59 | 0.48 | 0.083 |
| Spring Run Middle | | 0.43 | 0.49 | 1.14 | 0.63 | 0.273 |
| Spring Run Bottom | | 0.50 | 0.61 | 1.23 | 0.70 | 0.275 |
| Dumpling Run @Spring | 2006 | 0.12 | 0.16 | 0.36 | 0.19 | 0.088 |
| Dumpling Run Bottom | | 0.13 | 0.18 | 0.40 | 0.21 | 0.097 |
| Spring Run @Spring | | 0.34 | 0.41 | 0.55 | 0.41 | 0.079 |
| Spring Run Middle | | 0.39 | 0.41 | 0.57 | 0.45 | 0.074 |
| Spring Run Bottom | | 0.42 | 0.48 | 0.62 | 0.50 | 0.078 |
| Dumpling Run @Spring | 2007 | 0.00 | 0.16 | 0.22 | 0.14 | 0.077 |
| Dumpling Run Bottom | | 0.02 | 0.18 | 0.25 | 0.16 | 0.082 |
| Spring Run @Spring | | 0.02 | 0.39 | 0.41 | 0.33 | 0.154 |
| Spring Run Middle | | 0.00 | 0.51 | 0.69 | 0.45 | 0.234 |
| Spring Run Bottom | | 0.02 | 0.59 | 0.63 | 0.49 | 0.233 |
| Dumpling Run @Spring | 2008 | 0.12 | 0.15 | 0.26 | 0.17 | |
| Dumpling Run Bottom | | 0.13 | 0.16 | 0.31 | 0.19 | |
| Spring Run @Spring | | 0.36 | 0.42 | 0.58 | 0.44 | |
| Spring Run Middle | | 0.48 | 0.52 | 0.64 | 0.55 | |
| Spring Run Bottom | | 0.62 | 0.65 | 0.71 | 0.65 | |
| Nitrite-N (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 0.001 | 0.001 | 0.005 | 0.001 | 0.002 |
| Dumpling Run Bottom | | 0.001 | 0.001 | 0.014 | 0.003 | 0.006 |
| Spring Run @Spring | | 0.001 | 0.001 | 0.003 | 0.001 | 0.001 |

| | | | | | | |
|-----------------------|-------|-------|-------|-------|-------|-------|
| Spring Run Middle | | 0.001 | 0.007 | 0.023 | 0.009 | 0.009 |
| Spring Run Bottom | | 0.001 | 0.006 | 0.029 | 0.01 | 0.011 |
| Dumpling Run @Spring | 2006 | 0.001 | 0.001 | 0.001 | 0.001 | 0 |
| Dumpling Run Bottom | | 0.001 | 0.001 | 0.001 | 0.001 | 0 |
| Spring Run @Spring | | 0.001 | 0.001 | 0.001 | 0.001 | 0 |
| Spring Run Middle | | 0.004 | 0.009 | 0.031 | 0.014 | 0.01 |
| Spring Run Bottom | | 0.004 | 0.007 | 0.016 | 0.008 | 0.004 |
| Dumpling Run @Spring | | 2007 | 0.001 | 0.001 | 0.001 | 0.001 |
| Dumpling Run Bottom | 0.001 | | 0.001 | 0.001 | 0.001 | 0 |
| Spring Run @Spring | 0.001 | | 0.001 | 0.001 | 0.001 | 0 |
| Spring Run Middle | 0.001 | | 0.01 | 0.018 | 0.01 | 0.006 |
| Spring Run Bottom | 0.001 | | 0.003 | 0.006 | 0.003 | 0.003 |
| Dumpling Run @Spring | 2008 | 0.001 | 0.001 | 0.001 | 0.001 | |
| Dumpling Run Bottom | | 0.001 | 0.001 | 0.001 | 0.001 | |
| Spring Run @Spring | | 0.001 | 0.001 | 0.001 | 0.001 | |
| Spring Run Middle | | 0.001 | 0.013 | 0.031 | 0.014 | |
| Spring Run Bottom | | 0.001 | 0.001 | 0.018 | 0.007 | |
| TKN (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 0.041 | 0.115 | 0.758 | 0.24 | 0.273 |
| Dumpling Run Bottom | | 0.081 | 0.108 | 1.27 | 0.33 | 0.471 |
| Spring Run @Spring | | 0.099 | 0.15 | 1.89 | 0.44 | 0.711 |
| Spring Run Middle | | 0.214 | 0.305 | 1.46 | 0.51 | 0.475 |
| Spring Run Bottom | | 0.167 | 0.291 | 0.938 | 0.38 | 0.291 |
| Dumpling Run @Spring | 2006 | 0.035 | 0.203 | 0.233 | 0.163 | 0.079 |
| Dumpling Run Bottom | | 0.135 | 0.175 | 0.294 | 0.2 | 0.068 |
| Spring Run @Spring | | 0.091 | 0.214 | 0.287 | 0.201 | 0.066 |
| Spring Run Middle | | 0.181 | 0.214 | 1.09 | 0.387 | 0.355 |
| Spring Run Bottom | | 0.156 | 0.277 | 0.642 | 0.307 | 0.176 |
| Dumpling Run @Spring | 2007 | 0.117 | 0.167 | 0.196 | 0.161 | 0.035 |
| Dumpling Run Bottom | | 0.041 | 0.17 | 0.419 | 0.217 | 0.148 |
| Spring Run @Spring | | 0.089 | 0.149 | 0.243 | 0.155 | 0.051 |
| Spring Run Middle | | 0.167 | 0.288 | 0.527 | 0.302 | 0.122 |
| Spring Run Bottom | | 0.069 | 0.216 | 0.304 | 0.196 | 0.083 |
| Dumpling Run @Spring | 2008 | 0.009 | 0.109 | 0.233 | 0.111 | |
| Dumpling Run Bottom | | 0.009 | 0.083 | 0.341 | 0.150 | |
| Spring Run @Spring | | 0.009 | 0.131 | 0.271 | 0.145 | |
| Spring Run Middle | | 0.094 | 0.274 | 1.090 | 0.434 | |
| Spring Run Bottom | | 0.052 | 0.069 | 0.344 | 0.169 | |
| Total N (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 0.252 | 0.341 | 1.143 | 0.5 | 0.348 |
| Dumpling Run Bottom | | 0.274 | 0.364 | 1.784 | 0.63 | 0.583 |
| Spring Run @Spring | | 0.476 | 0.641 | 2.453 | 0.93 | 0.755 |
| Spring Run Middle | | 0.71 | 0.887 | 2.616 | 1.14 | 0.736 |
| Spring Run Bottom | | 0.688 | 0.877 | 2.197 | 1.09 | 0.574 |
| Dumpling Run @Spring | 2006 | 0.176 | 0.364 | 0.569 | 0.348 | 0.134 |
| Dumpling Run Bottom | | 0.296 | 0.356 | 0.695 | 0.407 | 0.146 |
| Spring Run @Spring | | 0.522 | 0.57 | 0.838 | 0.615 | 0.116 |
| Spring Run Middle | | 0.58 | 0.634 | 1.631 | 0.849 | 0.412 |
| Spring Run Bottom | | 0.601 | 0.734 | 1.216 | 0.818 | 0.227 |

| | | | | | | |
|----------------------|------|-------|-------|-------|-------|--------|
| Dumpling Run @Spring | 2007 | 0.197 | 0.312 | 0.405 | 0.305 | 0.073 |
| Dumpling Run Bottom | | 0.146 | 0.336 | 0.64 | 0.38 | 0.211 |
| Spring Run @Spring | | 0.165 | 0.517 | 0.644 | 0.485 | 0.167 |
| Spring Run Middle | | 0.306 | 0.775 | 1.01 | 0.766 | 0.258 |
| Spring Run Bottom | | 0.25 | 0.747 | 0.899 | 0.692 | 0.232 |
| Dumpling Run @Spring | 2008 | 0.160 | 0.259 | 0.370 | 0.280 | |
| Dumpling Run Bottom | | 0.170 | 0.274 | 0.652 | 0.341 | |
| Spring Run @Spring | | 0.370 | 0.640 | 0.694 | 0.581 | |
| Spring Run Middle | | 0.735 | 0.837 | 1.631 | 1.002 | |
| Spring Run Bottom | | 0.700 | 0.763 | 1.012 | 0.830 | |
| TP (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 0.007 | 0.028 | 0.059 | 0.028 | 0.019 |
| Dumpling Run Bottom | | 0.007 | 0.026 | 0.052 | 0.028 | 0.019 |
| Spring Run @Spring | | 0.013 | 0.025 | 0.046 | 0.028 | 0.014 |
| Spring Run Middle | | 0.049 | 0.075 | 0.166 | 0.086 | 0.046 |
| Spring Run Bottom | | 0.059 | 0.087 | 0.14 | 0.092 | 0.031 |
| Dumpling Run @Spring | 2006 | 0.022 | 0.054 | 0.124 | 0.06 | 0.035 |
| Dumpling Run Bottom | | 0.02 | 0.044 | 0.134 | 0.052 | 0.041 |
| Spring Run @Spring | | 0.026 | 0.049 | 0.261 | 0.08 | 0.09 |
| Spring Run Middle | | 0.081 | 0.103 | 0.179 | 0.122 | 0.044 |
| Spring Run Bottom | | 0.012 | 0.103 | 0.143 | 0.094 | 0.053 |
| Dumpling Run @Spring | 2007 | 0.033 | 0.041 | 0.046 | 0.039 | 0.005 |
| Dumpling Run Bottom | | 0.026 | 0.038 | 0.047 | 0.037 | 0.008 |
| Spring Run @Spring | | 0.033 | 0.036 | 0.068 | 0.043 | 0.014 |
| Spring Run Middle | | 0.072 | 0.106 | 0.293 | 0.128 | 0.083 |
| Spring Run Bottom | | 0.068 | 0.085 | 0.101 | 0.084 | 0.012 |
| Dumpling Run @Spring | 2008 | 0.013 | 0.042 | 0.049 | 0.036 | |
| Dumpling Run Bottom | | 0.039 | 0.042 | 0.049 | 0.042 | |
| Spring Run @Spring | | 0.023 | 0.033 | 0.046 | 0.034 | |
| Spring Run Middle | | 0.049 | 0.143 | 0.176 | 0.129 | |
| Spring Run Bottom | | 0.029 | 0.124 | 0.140 | 0.104 | |
| TSS (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 1.15 | 4.50 | 45.00 | 12.88 | 17.423 |
| Dumpling Run Bottom | | 1.15 | 2.08 | 43.00 | 9.58 | 16.588 |
| Spring Run @Spring | | 1.00 | 1.58 | 78.00 | 14.38 | 31.175 |
| Spring Run Middle | | 1.15 | 5.50 | 81.00 | 17.36 | 31.281 |
| Spring Run Bottom | | 4.00 | 6.50 | 72.00 | 18.50 | 26.629 |
| Dumpling Run @Spring | 2006 | 1.15 | 1.15 | 4.00 | 1.63 | 1.164 |
| Dumpling Run Bottom | | 3.00 | 5.50 | 8.00 | 5.50 | 2.258 |
| Spring Run @Spring | | 1.15 | 2.58 | 11.00 | 4.08 | 3.933 |
| Spring Run Middle | | 3.00 | 5.00 | 28.00 | 8.67 | 9.522 |
| Spring Run Bottom | | 3.00 | 7.00 | 19.00 | 8.50 | 5.431 |
| Dumpling Run @Spring | 2007 | 0.80 | 2.50 | 5.00 | 2.43 | 1.597 |
| Dumpling Run Bottom | | 0.80 | 4.50 | 7.00 | 3.97 | 2.246 |
| Spring Run @Spring | | 2.00 | 3.50 | 6.00 | 3.67 | 1.633 |
| Spring Run Middle | | 0.80 | 3.00 | 28.00 | 6.80 | 10.442 |
| Spring Run Bottom | | 3.00 | 6.00 | 7.00 | 5.83 | 1.472 |
| Dumpling Run @Spring | 2008 | 1.15 | 3.00 | 6.00 | 3.06 | |
| Dumpling Run Bottom | | 1.15 | 3.00 | 6.00 | 3.23 | |

| | | | | | | |
|------------------------|------|------|------|-------|-------|-------|
| Spring Run @Spring | | 1.15 | 1.15 | 11.00 | 3.69 | |
| Spring Run Middle | | 1.15 | 3.00 | 28.00 | 8.43 | |
| Spring Run Bottom | | 2.00 | 6.00 | 7.00 | 5.00 | |
| Turbidity (NTU) | | | | | | |
| Dumpling Run @Spring | 2005 | 0.45 | 0.9 | 22.95 | 7.52 | 10.6 |
| Dumpling Run Bottom | | 1.24 | 2.12 | 43.8 | 10.29 | 16.83 |
| Spring Run @Spring | | 1.03 | 1.95 | 18.42 | 5.22 | 6.84 |
| Spring Run Middle | | 1.31 | 3.4 | 36 | 9.58 | 13.52 |
| Spring Run Bottom | | 1.96 | 3.15 | 51.3 | 13.4 | 19.66 |
| Dumpling Run @Spring | 2006 | 0.67 | 0.97 | 2.15 | 1.14 | 0.59 |
| Dumpling Run Bottom | | 2.71 | 3.54 | 4.31 | 3.61 | 0.58 |
| Spring Run @Spring | | 3.14 | 4.78 | 7.47 | 4.86 | 1.66 |
| Spring Run Middle | | 2.43 | 3.65 | 12.8 | 5.27 | 4 |
| Spring Run Bottom | | 4.82 | 5.82 | 7.88 | 6.02 | 1.27 |
| Dumpling Run @Spring | 2007 | 1.2 | 1.58 | 6.82 | 2.49 | 2.18 |
| Dumpling Run Bottom | | 3.32 | 3.82 | 6.89 | 4.45 | 1.33 |
| Spring Run @Spring | | 2.27 | 4.98 | 7.27 | 4.88 | 1.72 |
| Spring Run Middle | | 1.95 | 3.11 | 25.1 | 7.43 | 9.04 |
| Spring Run Bottom | | 3.3 | 4.86 | 9.82 | 5.4 | 2.43 |
| Dumpling Run @Spring | 2008 | 1.07 | 1.28 | 1.94 | 1.39 | |
| Dumpling Run Bottom | | 2.49 | 3.50 | 4.92 | 3.68 | |
| Spring Run @Spring | | 2.27 | 2.59 | 7.47 | 3.54 | |
| Spring Run Middle | | 1.54 | 4.35 | 12.80 | 5.02 | |
| Spring Run Bottom | | 3.30 | 3.45 | 6.38 | 4.54 | |
| BOD5 (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 1.01 | 1.54 | 3.13 | 1.81 | 0.88 |
| Dumpling Run Bottom | | 1.18 | 1.52 | 2.68 | 1.63 | 0.55 |
| Spring Run @Spring | | 0.86 | 0.99 | 1.58 | 1.07 | 0.26 |
| Spring Run Middle | | 0.45 | 0.91 | 2.47 | 1.26 | 0.83 |
| Spring Run Bottom | | 0.66 | 1.01 | 1.97 | 1.15 | 0.49 |
| Dumpling Run @Spring | 2006 | 0.76 | 1.4 | 1.97 | 1.35 | 0.46 |
| Dumpling Run Bottom | | 0.76 | 1.1 | 1.59 | 1.12 | 0.34 |
| Spring Run @Spring | | 0.3 | 0.65 | 1.76 | 0.77 | 0.53 |
| Spring Run Middle | | 0.45 | 0.76 | 1.44 | 0.86 | 0.39 |
| Spring Run Bottom | | 0.23 | 0.43 | 1.36 | 0.57 | 0.43 |
| Dumpling Run @Spring | 2007 | 0.15 | 0.68 | 1.06 | 0.61 | 0.33 |
| Dumpling Run Bottom | | 0.23 | 0.61 | 1.21 | 0.61 | 0.37 |
| Spring Run @Spring | | 0.15 | 0.42 | 0.83 | 0.42 | 0.24 |
| Spring Run Middle | | 0.15 | 0.53 | 1.74 | 0.69 | 0.58 |
| Spring Run Bottom | | 0.15 | 0.42 | 1.29 | 0.53 | 0.41 |
| Dumpling Run @Spring | 2008 | 0.68 | 0.90 | 1.29 | 0.94 | |
| Dumpling Run Bottom | | 0.60 | 0.96 | 1.29 | 0.97 | |
| Spring Run @Spring | | 0.30 | 0.48 | 0.96 | 0.62 | |
| Spring Run Middle | | 0.30 | 0.71 | 1.44 | 0.76 | |
| Spring Run Bottom | | 0.10 | 0.75 | 1.16 | 0.65 | |
| DO (mg/L) | | | | | | |
| Dumpling Run @Spring | 2005 | 9.2 | 10.3 | 11.1 | 10.2 | 0.62 |
| Dumpling Run Bottom | | 9.4 | 10.4 | 11.5 | 10.5 | 0.69 |
| Spring Run @Spring | | 10 | 10.3 | 11.5 | 10.6 | 0.63 |

| | | | | | | |
|-----------------------------|------|------|-------|-------|-------|--------|
| Spring Run Middle | | 10.2 | 10.3 | 11.4 | 10.6 | 0.53 |
| Spring Run Bottom | | 10 | 10.6 | 11.2 | 10.6 | 0.48 |
| Dumpling Run @Spring | 2006 | 9.7 | 10.6 | 12.2 | 10.7 | 0.98 |
| Dumpling Run Bottom | | 9.8 | 10.6 | 12.2 | 10.7 | 0.86 |
| Spring Run @Spring | | 9.8 | 10.5 | 12.8 | 10.8 | 1.17 |
| Spring Run Middle | | 9.8 | 10.4 | 12.6 | 10.7 | 1.09 |
| Spring Run Bottom | | 9.9 | 10.7 | 12.6 | 10.9 | 1 |
| Dumpling Run @Spring | | 2007 | 10.2 | 10.5 | 11.2 | 10.6 |
| Dumpling Run Bottom | 9.7 | | 10.3 | 10.9 | 10.3 | 0.39 |
| Spring Run @Spring | 8.9 | | 10.1 | 11.2 | 10.2 | 0.82 |
| Spring Run Middle | 9.6 | | 10.3 | 10.9 | 10.3 | 0.49 |
| Spring Run Bottom | 9.9 | | 10.4 | 11.1 | 10.5 | 0.45 |
| Dumpling Run @Spring | 2008 | 9.1 | 10.0 | 11.2 | 10.1 | |
| Dumpling Run Bottom | | 9.5 | 10.4 | 10.9 | 10.3 | |
| Spring Run @Spring | | 9.7 | 10.1 | 10.4 | 10.0 | |
| Spring Run Middle | | 9.5 | 10.0 | 10.3 | 9.9 | |
| Spring Run Bottom | | 9.5 | 9.9 | 11.0 | 10.1 | |
| pH | | | | | | |
| Dumpling Run @Spring | 2005 | 7.4 | 7.8 | 8 | 7.7 | 0.24 |
| Dumpling Run Bottom | | 7.5 | 8.1 | 8.5 | 8 | 0.36 |
| Spring Run @Spring | | 7.4 | 7.9 | 8.2 | 7.8 | 0.31 |
| Spring Run Middle | | 7.3 | 7.8 | 8 | 7.8 | 0.25 |
| Spring Run Bottom | | 7.2 | 7.5 | 7.7 | 7.5 | 0.23 |
| Dumpling Run @Spring | 2006 | 7.3 | 7.6 | 7.8 | 7.6 | 0.17 |
| Dumpling Run Bottom | | 7.4 | 7.8 | 8.4 | 7.8 | 0.38 |
| Spring Run @Spring | | 7.3 | 7.7 | 8.2 | 7.7 | 0.31 |
| Spring Run Middle | | 7.3 | 7.7 | 8.4 | 7.7 | 0.42 |
| Spring Run Bottom | | 7.3 | 7.6 | 8.1 | 7.6 | 0.31 |
| Dumpling Run @Spring | 2007 | 7.1 | 7.5 | 7.9 | 7.5 | 0.34 |
| Dumpling Run Bottom | | 7.3 | 7.6 | 8.2 | 7.7 | 0.38 |
| Spring Run @Spring | | 7.2 | 7.7 | 8.2 | 7.7 | 0.33 |
| Spring Run Middle | | 6.9 | 7.5 | 8.1 | 7.5 | 0.44 |
| Spring Run Bottom | | 6.9 | 7.5 | 8.2 | 7.5 | 0.44 |
| Dumpling Run @Spring | 2008 | 7.1 | 7.3 | 7.8 | 7.4 | |
| Dumpling Run Bottom | | 7.4 | 8.0 | 8.3 | 7.8 | |
| Spring Run @Spring | | 7.3 | 7.8 | 8.2 | 7.8 | |
| Spring Run Middle | | 6.9 | 7.7 | 8.1 | 7.6 | |
| Spring Run Bottom | | 6.9 | 7.6 | 8.2 | 7.6 | |
| Conductivity (us/cm) | | | | | | |
| Dumpling Run @Spring | 2005 | 45.8 | 286.9 | 372 | 260 | 112.41 |
| Dumpling Run Bottom | | 48.1 | 283.5 | 352 | 257 | 106.74 |
| Spring Run @Spring | | 64.6 | 296.1 | 390 | 269 | 109.38 |
| Spring Run Middle | | 44.9 | 255 | 284 | 223 | 88.65 |
| Spring Run Bottom | | 45.1 | 247.5 | 276 | 213 | 85.87 |
| Dumpling Run @Spring | 2006 | 39 | 254.2 | 372 | 239.6 | 108.86 |
| Dumpling Run Bottom | | 44.9 | 263.9 | 352 | 241.3 | 103.48 |
| Spring Run @Spring | | 49 | 330.3 | 391.1 | 263 | 153.97 |
| Spring Run Middle | | 37.7 | 255.2 | 284 | 211.2 | 93.67 |
| Spring Run Bottom | | 36.3 | 253.2 | 276 | 213 | 90.6 |

| | | | | | | |
|----------------------|-------------|-------|-------|-------|-------|-------|
| Dumpling Run @Spring | 2007 | 234 | 257 | 345 | 271.7 | 39.83 |
| Dumpling Run Bottom | | 253 | 271.5 | 340 | 283.2 | 31.72 |
| Spring Run @Spring | | 271 | 358 | 375 | 347.5 | 38.3 |
| Spring Run Middle | | 241 | 253.5 | 277 | 257.2 | 14.26 |
| Spring Run Bottom | | 240 | 255 | 265 | 253.3 | 9.33 |
| Dumpling Run @Spring | 2008 | 234.0 | 247.0 | 356.0 | 266.6 | |
| Dumpling Run Bottom | | 253.0 | 266.0 | 355.0 | 280.0 | |
| Spring Run @Spring | | 288.0 | 378.0 | 390.0 | 363.4 | |
| Spring Run Middle | | 241.0 | 255.0 | 279.0 | 256.0 | |
| Spring Run Bottom | | 240.0 | 252.0 | 276.0 | 254.2 | |

Appendix 2. Laboratory Methods for Water Quality Parameters.

| Parameter | Method |
|-----------------------------|---------------|
| Ammonia Nitrogen | EPA 350.2 |
| Nitrate | EPA 353.2 |
| Nitrite | EPA 353.2 |
| * Ortho Phosphate | HACH 8048 |
| Total Phosphate | EPA 365.2 |
| Total Kjeldahl Nitrogen | EPA 351.2 |
| Total Suspended Solids | SM 2540D |
| * Turbidity | HACH 2100N |
| Biochemical Oxygen Demand 5 | SM5210B |

Appendix 3. WV Save Our Streams Macroinvertebrate Assessment July 2003

| Station 1 (catch-and-release) | Station 2 (catch-and-release) | Station 3 (above hatchery) |
|--------------------------------------|--------------------------------------|-----------------------------------|
| Ephemeroptera (mayflies) | Ephemeroptera (mayflies) | Ephemeroptera (mayflies) |
| <i>Baetidae</i> 73 | <i>Ephemerellidae</i> 1 | <i>Isonychiidae</i> 2 |
| <i>Heptageniidae</i> 2 | <i>Heptageniidae</i> 4 | <i>Ephemerellidae</i> 3 |
| Trichoptera (caddisflies) | <i>Baetidae</i> 45 | <i>Baetidae</i> 30 |
| <i>Rhyacophilidae</i> 2 | Plecoptera (stoneflies) | Plecoptera (stoneflies) |
| <i>Hydropsychidae</i> 13 | <i>Capniidae</i> 1 | <i>Capniidae</i> 17 |
| Diptera (true flies) | <i>Chloroperlidae</i> 1 | <i>Perlodidae</i> 6 |
| <i>Simuliidae</i> 8 | <i>Perlodidae</i> 4 | Trichoptera (caddisflies) |
| <i>Chironomidae</i> 67 | Trichoptera (caddisflies) | <i>Rhyacophilidae</i> 3 |
| Amphipoda (scuds) | <i>Glossosomatidae</i> 2 | <i>Hydropsychidae</i> 17 |
| <i>Gammaridae</i> 31 | <i>Rhyacophilidae</i> 1 | Coleoptera (beetles) |
| Total 196 | <i>Hydropsychidae</i> 18 | <i>Elmidae</i> 12 |
| | Coleoptera (beetles) | <i>Psephenidae</i> 1 |
| | <i>Elmidae</i> 4 | Diptera (true flies) |
| | Diptera (true flies) | <i>Dixidae</i> 1 |
| | <i>Simuliidae</i> 16 | <i>Simuliidae</i> 12 |
| | <i>Chironomidae</i> 37 | <i>Chironomidae</i> 6 |
| | Amphipoda (scuds) | Amphipoda (scuds) |
| | <i>Gammaridae</i> 125 | <i>Gammaridae</i> 60 |
| | Total 259 | Total 170 |

Appendix 4. Assessing the Condition of the Macroinvertebrate Communities of Spring Run
(Tim Craddock, Citizen's Monitoring Coordinator).

| Integrity ratings for stations through study period | | | | | | | |
|---|------|----------|------------|---------|----------|-------|--|
| Study periods | Poor | Marginal | Suboptimal | Optimal | Averages | RPD | Stations |
| 2005 - 2006 | 7 | 9 | 4 | 0 | 49.5 | 30.63 | All stations |
| 2007 - 2008 | 0 | 5 | 14 | 1 | 67.4 | | |
| 2005 - 2006 | 6 | 4 | 2 | 0 | 47.0 | 32.61 | All Spring Run stations |
| 2007 - 2008 | 0 | 4 | 8 | 0 | 65.3 | | |
| 2005 - 2006 | 1 | 5 | 2 | 0 | 53.3 | 27.93 | Dumpling Run stations |
| 2007 - 2008 | 0 | 1 | 6 | 1 | 70.5 | | |
| 2005 - 2006 | 4 | 4 | 0 | 0 | 43.3 | 34.20 | Spring Run stations downstream of hatchery |
| 2007 - 2008 | 0 | 4 | 4 | 0 | 61.2 | | |

| Station Codes | May-05 | Oct-05 | Jun-06 | Oct-06 |
|--------------------|---------|----------|---------|----------|
| Spring Run (0.4) | SR1-505 | SR1-1005 | SR1-606 | SR1-1006 |
| Spring Run (1.6) | SR2-505 | SR2-1005 | SR2-606 | SR2-1006 |
| Spring Run (2.3) | SR3-505 | SR3-1005 | SR3-606 | SR3-1006 |
| Dumpling Run (1.4) | DR1-505 | DR1-1005 | DR1-606 | DR1-1006 |
| Dumpling Run (2.2) | DR2-505 | DR2-1005 | DR2-606 | DR2-1006 |

| Station Codes | May-07 | Oct-07 | Apr-08 | Oct-08 |
|--------------------|----------|----------|----------|----------|
| Spring Run (0.4) | SR1-0507 | SR1-1007 | SR1-0408 | SR1-1008 |
| Spring Run (1.6) | SR2-0507 | SR2-1007 | SR2-0408 | SR2-1008 |
| Spring Run (2.3) | SR3-0507 | SR3-1007 | SR3-0408 | SR3-1008 |
| Dumpling Run (1.4) | DR1-0507 | DR1-1007 | DR1-0408 | DR1-1008 |
| Dumpling Run (2.2) | DR2-0507 | DR2-1007 | DR2-0408 | DR2-1008 |

| Codes | Total Taxa | EPT Taxa | Biotic Index | % EPT | % Dominant | % Tolerant | Stream Index | Integrity |
|----------|------------|----------|--------------|-------|------------|------------|--------------|------------|
| SR1-0505 | 11 | 5 | 5.39 | 8.9 | 65.5 | 19.1 | 48.7 | Marginal |
| SR2-0505 | 14 | 7 | 5.76 | 38.5 | 42.3 | 54.0 | 58.8 | Marginal |
| SR3-0505 | 16 | 10 | 4.78 | 40.4 | 41.3 | 16.1 | 70.3 | Suboptimal |
| DR1-0505 | 16 | 9 | 4.80 | 31.0 | 43.8 | 16.7 | 68.2 | Suboptimal |
| DR2-0505 | 10 | 5 | 4.78 | 14.7 | 80.4 | 3.1 | 50.1 | Marginal |
| SR1-1005 | 6 | 3 | 5.01 | 1.6 | 77.6 | 4.7 | 41.7 | Poor |
| SR2-1005 | 11 | 5 | 7.20 | 4.6 | 76.9 | 78.5 | 31.1 | Poor |
| SR3-1005 | 8 | 4 | 4.89 | 6.4 | 87.2 | 2.7 | 44.1 | Poor |

| Codes | Total Taxa | EPT Taxa | Biotic Index | % EPT | % Dominant | % Tolerant | Stream Index | Integrity |
|----------|------------|----------|--------------|-------|------------|------------|--------------|------------|
| DR1-1005 | 14 | 9 | 4.54 | 25.3 | 68.8 | 0.8 | 63.5 | Suboptimal |
| DR2-1005 | 11 | 7 | 4.85 | 7.1 | 87.9 | 2.5 | 50.4 | Marginal |
| SR1-0606 | 9 | 6 | 5.03 | 13.7 | 53.1 | 15.0 | 53.4 | Marginal |
| SR2-0606 | 12 | 5 | 6.55 | 23.9 | 64.3 | 64.8 | 42.0 | Poor |
| SR3-0606 | 12 | 8 | 4.62 | 32.8 | 55.9 | 7.8 | 63.5 | Suboptimal |
| DR1-0606 | 7 | 2 | 5.00 | 1.0 | 87.6 | 1.9 | 39.5 | Poor |
| DR2-0606 | 10 | 7 | 4.82 | 10.4 | 82.2 | 4.3 | 51.2 | Marginal |
| SR1-1006 | 9 | 4 | 5.32 | 7.0 | 68.1 | 14.6 | 45.9 | Marginal |
| SR2-1006 | 10 | 2 | 7.31 | 3.3 | 80.5 | 80.9 | 24.8 | Poor |
| SR3-1006 | 7 | 2 | 4.90 | 2.2 | 90.2 | 0.4 | 39.4 | Poor |
| DR1-1006 | 12 | 6 | 4.78 | 10.2 | 83.6 | 1.8 | 51.6 | Marginal |
| DR2-1006 | 11 | 7 | 4.85 | 10.2 | 83.4 | 4.9 | 51.5 | Marginal |
| SR1-0507 | 11 | 6 | 4.62 | 38.8 | 45.3 | 9.3 | 62.8 | Suboptimal |
| SR2-0507 | 9 | 5 | 5.04 | 45.1 | 34.4 | 20.0 | 60.7 | Suboptimal |
| SR3-0507 | 14 | 9 | 4.04 | 71.0 | 34.4 | 9.0 | 78.6 | Suboptimal |
| DR1-0507 | 14 | 10 | 3.88 | 53.2 | 40.3 | 3.9 | 76.6 | Suboptimal |
| DR2-0507 | 12 | 7 | 4.49 | 44.9 | 51.3 | 1.7 | 66.3 | Suboptimal |
| SR1-1007 | 13 | 7 | 4.66 | 63.0 | 34.2 | 9.6 | 73.2 | Suboptimal |
| SR2-1007 | 11 | 4 | 5.92 | 25.5 | 34.8 | 35.5 | 53.3 | Marginal |
| SR3-1007 | 16 | 10 | 4.23 | 47.4 | 43.2 | 0.9 | 76.8 | Suboptimal |
| DR1-1007 | 21 | 12 | 4.04 | 59.4 | 8.7 | 27.0 | 88.2 | Optimal |
| DR2-1007 | 16 | 10 | 4.63 | 29.8 | 50.8 | 8.7 | 69.7 | Suboptimal |
| SR1-0408 | 13 | 8 | 5.07 | 17.9 | 66.9 | 10.7 | 57.3 | Marginal |
| SR2-0408 | 11 | 6 | 5.79 | 23.6 | 38.2 | 38.2 | 53.8 | Marginal |
| SR3-0408 | 11 | 10 | 4.72 | 28.6 | 59.1 | 9.5 | 66.7 | Suboptimal |
| DR1-0408 | 14 | 9 | 4.36 | 31.3 | 59.3 | 5.1 | 66.0 | Suboptimal |
| DR2-0408 | 13 | 8 | 4.83 | 25.2 | 49.5 | 18.8 | 61.4 | Suboptimal |
| SR1-1008 | 13 | 8 | 4.74 | 38.2 | 27.3 | 13.2 | 69.6 | Suboptimal |
| SR2-1008 | 8 | 4 | 5.26 | 40.1 | 28.8 | 18.5 | 58.6 | Marginal |
| | | | | | | | | |

| Codes | Total Taxa | EPT Taxa | Biotic Index | % EPT | % Dominant | % Tolerant | Stream Index | Integrity |
|----------|------------|----------|--------------|-------|------------|------------|--------------|------------|
| SR3-1008 | 16 | 10 | 4.64 | 36.4 | 43.6 | 7.6 | 71.9 | Suboptimal |
| DR1-1008 | 15 | 9 | 3.88 | 51.4 | 37.5 | 2.4 | 76.6 | Suboptimal |
| DR2-1008 | 13 | 9 | 4.52 | 16.6 | 74.6 | 1.5 | 59.5 | Marginal |